

LA-UR 93-0189

LA-UR--93-189

DE93 007341

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

RECEIVED
FEB 11 1993
OSTI

TITLE DNA Repair and Radiation Sensitivity in Mammalian Cells

AUTHOR(S) David Jen-Chi Chen, Murray A. Stackhouse, and Davis Shao-Shuan Chen

SUBMITTED TO Proceeding for the International Workshop on Radiation Effects
Taipei, Taiwan
October 18, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or approval by the United States Government or any agency thereof. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

DNA repair and Radiation Sensitivity in Mammalian Cells

**David Jen-Chi Chen, Murray Stackhouse and
Davis Shao-shuan Chen⁺**

**Life Sciences Division, Los Alamos National Laboratory, Los Alamos, New
Mexico and ⁺Department of Radiation Oncology, School of Medicine,
University of Rochester, Rochester, New York**

**Proceeding for the International Workshop on Radiation Effects, Taipei,
Taiwan, October 18, 1992**

Abstract

Ionizing radiation induces various types of damage in mammalian cells including DNA single-strand breaks, DNA double-strand breaks (DSB), DNA-protein cross links, and altered DNA bases. Although human cells can repair many of these lesions there is little detailed knowledge of the nature of the genes and the encoded enzymes that control these repair processes. We report here on the cellular and genetic analyses of DNA double-strand break repair deficient mammalian cells. It has been well established that the DNA double-strand break is one of the major lesions induced by ionizing radiation. Utilizing rodent repair-deficient mutant, we have shown that the gene(s) responsible for DNA double-strand break repair is also responsible for the cellular expression of radiation sensitivity. The molecular genetic analysis of DSB repair in rodent/human hybrid cells indicate that at least 6 different genes in mammalian cells are responsible for the repair of radiation-induced DNA double-strand breaks. Mapping and the prospect of cloning of human radiation repair genes are reviewed.

There is evidence that DNA repair capacities differ among individuals, and there are well known instances of inherited susceptibility to cancer. Our understanding of the genetics of mammalian DNA repair suggest that a highly complex system is involved. Therefore, the likelihood that there are a number of alleles in the human population that affect susceptibility to radiation effects when present in the heterozygous state must be taken very seriously. Understanding the molecular and genetic basis of radiation sensitivity and DNA repair in man will provide a rational foundation to predict the individual risk associated with radiation exposure and to prevent radiation-induced genetic damage in the human population.

Introduction

A number of laboratories have investigated the mechanisms by which cell killing, mutation, and neoplastic transformation are induced in rodent or human cells in vitro following exposure to UV light, X-irradiation and a variety of chemical carcinogens. When such cells are allowed to repair DNA prior to cell division, repair of DNA damage has been linked to the biological recovery of cells under these conditions (1-3). These findings support the hypothesis that the response of eukaryotes to DNA damaging agents is determined by the effectiveness of a variety of DNA repair systems. Therefore, understanding the nature of the repair and misrepair processes is central to the elucidation of the mechanisms through which subsequent adverse health effects such as mutations, fetal malformations, and cancers are expressed.

Ionizing radiation produces a variety of damage in DNA including DNA-protein cross-links, DNA single strand breaks (SSB), DNA double strand breaks (DSB), and base damage. In prokaryotes and lower eukaryotes, a number of genes involved in the repair of DNA damage caused by ionizing radiation have been identified. In some cases these genes have been cloned and their role in different repair pathways partially characterized (4). These studies have been considerably helped by the use of mutant strains. However, DNA repair mechanisms in mammalian cells are less well understood, primarily due to the lack of suitable mutant cell lines. However, similar approaches to those used in prokaryotes and lower eukaryotes are now possible in mammalian cells with the advent of recombinant DNA techniques and the isolation of a number of mammalian cell mutants that are defective in DNA repair.

In this report we will review recent studies on the genetics of DNA repair in mammalian cells using mutants that are deficient in the repair of radiation-induced DNA double-strand breaks.

Human genetic syndromes

Mutants of microorganisms sensitive to various DNA damaging agents have been of immense importance in helping to elucidate the major pathways of DNA repair in bacteria. In yeast, more than 50 mutants have been isolated that express sensitivity to UV, X-rays or chemical carcinogens and are shown to be deficient in certain repair genes. In humans, individuals afflicted by certain autosomal recessive syndromes are at high risk for cancer. Cells from these patients exhibit both chromosome instability and increased sensitivity to specific DNA damaging agents in culture. Cleaver (5,6) has provided evidence that fibroblasts from

xeroderma pigmentosum (XP) patients are defective in their ability to repair specific lesions formed in DNA as a result of exposure to UV light. Increased sensitivity to X-Irradiation of cultured fibroblasts has been found in a number of autosomal recessive diseases, such as: ataxia telangiectasia (AT) (7), Fanconi's anemia (FA) (8), Cockayne's syndrome (CS) (9), Gardner syndrome (GS)(9), and Rothmund Thomson syndrome (RTS) (10). These genetically disordered patients are apparently deficient in some form of DNA repair. Smith and Paterson (10) have suggested that the primary defect of RTS may be an aberrant gene control function which results in a misdirected repression of a DNA repair gene(s). Several of the XP and CS genes have been identified recently (11-13). However, most of these genes or gene products that are affected in all of these cancer prone disorders are unknown.

Genetic analysis of radiation-sensitive mutants

As an approach to the elucidation and the understanding of mammalian DNA repair, a number of Chinese hamster ovary (CHO) cell lines that are sensitive to UV radiation or chemical carcinogens have been isolated (14-16). A number of ionizing radiation sensitive rodent mutants have also been isolated (14-19). Thirteen of these mutants have been shown to be deficient in rejoining of radiation-induced DNA DSB based on results obtained by neutral elution analysis or pulsed field gel electrophoresis (20-23). Genetic complementation analysis of different mutant cell lines with the same phenotype is often used to determine the number of genetic loci involved in the expression of a particular trait. As indicated in Table 1, a total of 13 rodent radiation sensitive mutants have been identified as deficient in DNA DSB repair. Mutants within the same category belong to the same complementation group. Most of the repair deficient mutant cell lines are complemented by normal human fibroblasts or lymphocytes. The first four groups of mutants belong to different complementation groups. All the mouse DSB repair-deficient mouse mutants belong to different complementation groups. Therefore, up to 10 human genes are responsible for the repair of ionizing radiation-induced DNA DSBs.

Cellular repair capability of DSB repair mutant cells

One of the DSB rejoining deficient mutants, XRS-6, has been characterized extensively. XRS-6 cells are highly sensitive to X-rays in terms of cell killing (19), chromosome aberrations (24,25), enhanced mutation induction (26,27), G2 block (28), and the enhanced radiosensitivity of DNA synthesis (24). Our results also indicate that XRS-

6 cells have no potential lethal damage repair (30), no sublethal damage repair (31), and no dose-rate effects (4). Biochemical analyses have also indicated that the ligase (32), topoisomerase (33), and AP endonuclease activities (unpublished results) of XRS-6 cells are normal and the only biochemical defect has been the inability to rejoin DNA DSBs (22,23). A summary of these studies is presented here.

<u>cell line</u>	<u>Origin</u>	<u>Complementation by human cells</u>	<u>Designed human gene</u>	<u>References</u>
1 XR-1 M10	CHO L5178Y	yes	XRCC4	23, 18
2 XRS-6 XRV15B	CHO V79	yes	XRCC5	29, 16
3 V3	CHO	yes		62
4 XR-V9B	V79	yes		63
5 BLM-2	CHO	yes		64
6 SL3-1478	Mouse L	yes		65
7 LX830	L5178Y	in progress		14
8 L5178Y/S L5178Y/SS	L5178Y L5178Y	in progress		21, 66
9 SX9	Mouse SR-1	in progress		67
10 SX10	Mouse SR-1	in progress		61

Table 1. Rodent DNA double-strand break repair-deficient mutants

Repair of radiation-induced DNA SSB and DSB

The yield of DNA SSB in CHO-K1 and XRS-6 cells upon exposure to increasing radiation dose was found to be the same (data not shown). Closure of DNA SSB proceeded at a similar rate in both CHO-K1 and XRS-6 cells with an initial SSB closure half-time of 3 min at 37°C (unpublished results). Induction of DNA DSB was estimated in these two cell lines by pulsed-field gel electrophoresis. Exposure to increasing radiation doses caused an increasing fraction of cell DNA to enter the agarose gel in both CHO-K1 and xrs-6 cells (data not shown). It has been assumed that an

increase in the fraction of DNA radioactivity entering the agarose gel reflected a decrease in the length of nuclear DNA and thus an increase in the frequency of DNA DSB. Rejoining of DNA DSB has been estimated by comparing the amount of DNA in the gel and the plug. After a dose of 50 Gy and a repair time of 4 h the xrs-6 cells showed a qualitatively reduced ability to rejoin breaks measured by PFGE compare to the parental cells. In parental, CHO-OK1 cells the radiation-induced dsbs were rapidly removed. However, the DNA breaks were closed in xrs-6 cells somewhat more slowly, with 50% of the breaks being closed in CHO-K1 and xrs-6 cells after 5 and 12 min respectively (33).

Effect of delayed plating in synchronized G1 cells

Both of the parental and mutant cells were synchronized in G1 by isoleucine depletion (34). These cultures were used for the delayed-plating PLD repair determination and for split-dose recovery in G1 cells. CHO-K1 and XRS-6 cells were inoculated into T-25 flasks and synchronized by isoleucine depletion. Estimated by autoradiographic analysis of cultures with a 60-min pulse of [³H]thymidine (10mCi/ml at a specific activity of 50 Ci/mmol), it was shown that between 10 and 12% of the cells were in S phase. The survival fraction of G1 phase-arrested CHO-K1 cells increased during postirradiation incubation in IL- medium. As described by Nagasawa et al., (), CHO-K1 cells displayed the typical pattern of PLD repair that saturated about 8 h postirradiation incubation. When PLD repair was measured at two or three different survival levels the recovery ratio increased proportionately with dose. However, XRS-6 mutant cells showed no PLD repair during the postirradiation holding period in IL-medium. Lack of PLD repair has also been observed in other DSB repair-deficient mutants.

Effects of plit-dose irradiation in synchronized G1 cells

We measured the split-dose recovery in parental and mutant cells by delivering two equal dose fractions at various time intervals to cells synchronized by isoleucine depletion (30). There is a large increase in survival for CHO-K1 cells as a function of the time between doses. The repair half-time for the parental cell line was about 90 min and an 4 - 5 fold increase in survival was seen when the doses were separated by 4 -12 hours. By contrast, there is no increase in survival for the sensitive mutant, XRS-6 cells during the interval of dose fraction.

Effects of low dose rate

Dose rate effects were measured in synchronized G1 cells. CHO K1 cells exhibited a large dose rate effect when survival curve generated from exposure to g-rays at 75 cGy/min and 15.3 cGy/h were compared (30). On the other hand, there was no apparent dose-rate effect for the X-ray-sensitive XRS-6 mutant cells, even when the cells were exposed to g-ray at the very low dose rate of 2.7 cGy/h (30).

Radiation mutagenesis

It has been suggested that radiation-induced deletion mutation may result from DNA double-strand breaks since experiments show that both chromosomal aberrations and mutations can be induced by treating cells with restriction endonucleases (35). Evans et al. (36) suggested that the repair deficiency could be responsible for the occurrence of residual DNA multilocus lesions following X-irradiation in the radiosensitive strain LY-S of the mouse lymphoma cell line (L5178Y) which results in its increased mutational response. In addition to mouse LY-S cells, enhanced radiation-induced mutation-induction (hypermutability) observed in several other DSB repair deficient mutants (37,38) suggests that the mutational events outnumber the lethal events induced by DSBs. A plot of mutation frequency versus survival have been analyzed in mutants M10 and XRS-6 (37,39). It is reasonably fitted by a common line suggesting a common lesion type leading to expressed mutations and cell killing in both these cell lines. These results suggest the likely involvement of DSB in mutation induction as well as cell killing in irradiated mammalian cells. It also seems likely that the high mutability of the DSB repair-deficient mutants result from its inability to remove dsb from the DNA resulting in high levels of deletions (40,41).

Cloning of human DSB repair genes

Present attempts to clone human DNA-repair genes rely on schemes which utilize human genomic DNA to complement rodent lines with defined deficiencies in DNA repair mechanisms. Once a complementing human gene can be identified within a deficient rodent background (introduced by DNA transfection), recombinant DNA technologies can be utilized to isolate and clone the human gene. In this approach, DNA from normal human cells is transfected into mutant hamster cells using the calcium phosphate precipitation method. Selective pressure is then applied to the transfected population by treatment with an appropriate agent (e.g., UV, X-rays, chemicals) under conditions that allow only transfected cells that have a normal or wild-type phenotype to survive. It is presumed that

resistant rodent clones have acquired the repair gene of interest by integration of the foreign human DNA. The repair-proficient transformants are often screened with human repetitive sequences to verify the presence of DNA of human origin within the transformants. The human Alu-family sequences constitute a built-in physical genetic marker that allow for the detection and localization of human DNA sequences in transformed hamster cells. In the past 10 years, 6 human DNA repair genes have been isolated. Westerveld et al., (42) Weber et al.,(43), Mudgett et al., (44) and Weeda et al, (45) have used similar strategies to clone 4 human repair genes utilizing CHO mutants deficient in DNA excision repair (ERCC1, ERCC2, ERCC3, ERCC5, ERCC6). A human repair gene responsible for the rejoining of DNA single strand breaks (XRCC 1) has also been cloned by Thompson et al. (46).

There are several major reasons for such slow progress in isolating human radiation DNA repair genes: 1) Current methods for cloning are dependent on functional complementation rather than use of molecular probes; 2) Mammalian cells in general are inefficient at taking up large DNA molecules. In the case of DSB repair-deficient mutants, such as XRS-6, the transfection frequency by cosmids is at least 10-50 fold less than normal CHO cells (unpublished results); 3) DSB repair-deficient mutants are sensitive to ionizing radiation, but radiation itself is not an ideal selection agent for the isolation of repair-proficient transformants.

To avoid all the possible phenotypic drawbacks of the repair-deficient mutants described above, advanced somatic and molecular genetic technologies have been suggested for the cloning of human DSB repair genes. It is well established that functional complementation can be achieved by somatic cell hybridization at a very high efficiency. The human chromosome that complements defective rodent repair phenotypes can be identified from the resulting hybrids between normal human cells and rodent mutants (477-49). Recent development of a high efficiency ($>1 \times 10^{-5}$) chromosome transfer technique provides an approach for construction of hybrids containing only a single specific human chromosome (50-51). If the radiation-sensitivity of rodent repair-deficient mutant cells can be complemented by a single human chromosome, it means that a repair gene(s) is located on this specific human chromosome. Somatic cell genetic techniques have also been developed to reduce the amount of human chromosomes present in a hybrid cell (52) Therefore, it is now possible to construct a human/mutant hybrid, which contains only a fragment of a specific human chromosome that can functionally complement the mutant cells. Subsequently, unique human sequences or transcripts located on a fragment of a human chromosome will be amplified and isolated by PCR technique using human specific Alu as the primer (53). These isolated human probes from a

known region of a specific human chromosome will then be used to clone the repair gene(s) in YACs which can accommodate large sizes of DNA (100-800 kb).

Mapping of human DSB repair genes

Identification of human chromosomes carrying genes that complement DNA repair deficiencies in rodent mutant cells has been done by somatic cell genetic techniques. Human cells were fused to rodent mutant cells and hybrids were selected at levels of DNA-damaging agent at which the mutant cells were hypersensitive. Concordance of hybrid clone resistance to the selecting agent with a particular human chromosome (or part of a human chromosome) identifies the location of the complementing gene -- Excision Repair Cross Complementing (ERCC) or X-ray Repair Cross Complementing (XRCC). Such studies have mapped ERCC1, ERCC2 and XRCC1 onto human chromosome 19q13.1-q13.3 (54-56), ERCC3 and ERCC5 onto chromosome 2 and 13, respectively (49), and XRCC4 onto chromosome 5 (50). Furthermore, similar preliminary data suggest the location of ERCC4 on chromosome 16p13.13-p13.3 (58), and that of XRCC2 and XRCC3 on chromosomes 7 and 14, respectively (59).

Radiation resistance or sensitivity of somatic cell hybrids constructed from the fusion of XRS-6 cells with primary human fibroblasts strongly correlated with the retention of human chromosome 2 isozyme and molecular markers. Disconcordanities between some chromosome 2 markers and the radiation resistance phenotype in some of the hybrid cells suggested the location of X-ray Repair Cross Complementing 5 (XRCC5) gene on the chromosome 2 (60). Introduction of human chromosome 2 by microcell-mediated chromosome transfer into the radiation sensitive XRS-6 cells resulted in hybrid cells in which the radiation sensitivity was complemented (39, 60).

DNA repair and radiation protection

There is evidence that DNA repair capacities differ among individuals, and there are well known instances of inherited susceptibility to cancer. Our understanding of the genetics of mammalian DNA repair suggest the involvement of a highly complex system. It also clear that certain DNA-repair-related genetic disorders are associated with autosomal recessive diseases, such as XP and AT. Even though human genetic disorders related to DNA DSB repair have not been identified yet, it might be expected the heterozygotic state of some of these DSB repair genes or these disease-associated alleles would have some phenotypic effect. This could have

quite an important effect on the population. Nagasawa et al., have measured radiation sensitivity among ten apparent normal human diploid fibroblast strains based on three different end points (26). They have found that three of these ten strains showed a moderate degree of hypersensitivity to X-rays by all three assays, falling within the range previously reported for AT heterozygotes (27). Therefore, the likelihood that there are a number of alleles in the human population that affect susceptibility to radiation effects when present in the heterozygous state must be taken very seriously. Understanding the molecular and genetic basis of radiation sensitivity and DNA repair in man will provide a rational foundation to predict the individual risk associated with radiation exposure and to prevent radiation-induced genetic damage in the human population.

Acknowledgment

We are indebted to Ms. Thu Nguyen and Jeanelle Ortize for technical assistance. This work was supported by the U.S. Department of Energy and National Institute of Health.

References

1. R.F. Jostes, Jr. and R.B. Painter, Evidence for repair of premutational damage from split x-ray doses, *Radiat. Res.* 84:586-590 (1980)
2. M. Terzaghi and J.B. Little, X-radiation-induced transformation in a C3H mouse embryo-derived cell line, *Cancer Res.* 36:1367-1374 (1976).
3. V. M. Maher, D. J. Dorney, A. L. Mendrala, B. Konze-Thomas and J. J. McCormick, DNA excision-repair processes in human cells can eliminate the cytotoxic and mutagenic consequences of ultraviolet irradiation, *Mutat. Res.* 62:311-323 (1979).
4. E. C. Friedberg, Deoxyribonucleic acid repair in the yeast *Saccharomyces cerevisiae*, *Microbiological Review* 52:70-102 (1988).
5. J. E. Cleaver, Defective repair replication of DNA in xeroderma pigmentosum, *Nature* 218:652-56 (1968).
6. J.E. Cleaver, Defective repair replication of DNA in xeroderma pigmentosum, *Nature* 218:652-56 (1968).
7. M.C. Paterson, and P.J. Smith, Ataxia telangiectasia: An inherited human disorder involving hypersensitivity to ionizing radiation and related DNA-damaging chemicals, *Ann. Rev. Genet.* 13:291-318 (1979).
8. J. E. Cleaver, DNA repair processes and their impairment in some human diseases. In: "Progress in Genetic Toxicology," D. Scott, B.A. Bridges, F. H. Sobels, eds. (Elsevier), pp. 29-42 (1977).
9. C.F. Arlett, and S.A. Harcourt, Survey of radiosensitivity in a variety of human cell strains, *Cancer Res.* 40:926-932 (1980).
10. P.J. Smith and M.C. Paterson, Enhanced radiosensitivity and defective repair in cultured fibroblasts derived from Rothmund Thomson syndrome patients, *Mutat. Res.* 94:213-228 (1982).
11. M. Bankmann, L. Prakash, S. Prakash, Yeast RAD 14 an human Xeroderma-Pigmentosum group A DNA repair genes encode homologous proteins, *Nature*

- 355:6360 (1992)
12. K. Tanaka, N. Miura, I. Satokata, I. Miyamoto, M. C. Yoshida, Y. Saioh, S. Kondo, A. Yasui, H. Okayama, Y. Okada, Analysis of a human DNA excision repair gene involved in group A Xeroderma-Pigmentosum and containing a Zinc-finger domain, *Nature* 348:6296 (1990).
 13. G. Weeda, R. C. A. Van Ham, W. Vermeulen, D. Bootsma, A. J. van der Eb, and J. H. J. Hoeijmakers, A presumed DNA helicase encoded by ERCC-3 is involved in the human repair disorders Xeroderma Pigmentosum and Cockayne's Syndrome, *Cell* 62:777-691 (1990).
 14. K. Sato, N. Hieda-Shiomi and J. Hama-Inaba, X-ray-sensitive mutant mouse cell with various sensitivities to chemical mutagens, *Mutat. Res.* 121:281-285 (1983).
 15. L. H. Thompson, DNA repair mutants, p.641-667, In M.M. Gottesman (ed), *Mole. Cell. Genet.* John Wiley & Sons, N.Y. (1985).
 16. M. Z. Zdzienicka and J.W.I.N. Simons, Mutagen-sensitive cell lines are obtained with a high frequency in V79 Chinese hamster cells, *Mutat. Res.* 178:235-244 (1987).
 17. L. F. Fuller, and R. B. Painter, A Chinese hamster ovary cell line hypersensitive to ionizing radiation and deficient in repair replication, *Mutat Res* 193:109-121 (1988).
 18. K. Sato and N. Hieda, Isolation and characterization of a mutant mouse lymphoma cell sensitive to methanesulfonate and X-rays, *Radiat. Res.* 78:167-171 (1979)
 19. P.A. Jeggo, X-ray sensitive mutants of Chinese hamster ovary cell line: radio-sensitive of DNA synthesis, *Mutat. Res.* 145:171-76 (1985)
 20. K. Eguchi-Kasai, T. Kosaka, K. Sato and I. Daneko, Reparability of DNA double-strand breaks and radiation sensitivity in five mammalian cell lines, *Int. J. Radiat. Biol.*, 59:97104 (1991)
 21. Evans, H. H., M. Ricanati, and M. Horng, Deficiency in DNA repair in mouse lymphoma strain L5178Y-S, *Proc. Natl. Acad. Sci. USA* 84:7562-7566 (1987)
 22. 28. Kemp, L. M., S. G. Sedgwick, and P. A. Jeggo, X-ray sensitive mutants of Chinese hamster ovary cells defective in double-strand break rejoining, *mutat. Res.* 132:189-196 (1984)
 23. Stamato, T. D. R. Weinstein, A. Giaccia and L. Mackenzie, Isolation of cell cycle-dependent gamma ray-sensitive Chinese hamster ovary cells, *Somat. Cell Genet.* 9:165-73 (1983)
 24. F. Darroudi and A. T. Natarajan, Cytogenetical characterization of Chinese hamster ovary X-ray-sensitive mutant cells xrs 5 and xrs 6, *Mutat. Res.* 212:123-135 (1989)
 25. T. A. K. Mussa, B. Singh and P. E. Bryant, Enhance mutability at the tk locus in the radiosensitive double-strand break repair mutant xrs5, *Mutat. Res.* 231:187-193 (1990)
 26. H. Nagasawa and J. Little, Radiosensitivities of ten apparently normal human diploid fibroblast strains to cell killing, G2-phase chromosomal aberrations, and cell cycle delay, *Cancer Res.* 48:4543-4538 (1988)
 27. H. Kraemer, Y. Shiloh and J. B. Little, Detection of ataxia telangiectasia heterozygous cell lines by post-irradiation cumulative labeling index: measurement with coded samples, *Cancer Res.* 47:398-402 (1987)
 28. K. F. Weibezahn, H. Lohrer and P. Herlich, Double-strand break repair and G2 block in Chinese hamster ovary cells and their radiosensitive mutants, *Mutat. Res.* 145:177-183 (1985)
 29. P. Jeggo, P.A. X-ray sensitive mutants of Chinese hamster ovary cell line: radio-sensitive of DNA synthesis, *Mutat. Res.* 145:171-76 (1985)
 30. H. Nagasawa, D.J. Chen and G.F. Strniste, Response of X-ray sensitive CHO cells to X-irradiation: 1. Effects of low-dose rates and the process of PLD repair in G1-phase, *Radiat. Res.* 118:559 -567 (1989)

31. D. J. Chen, G. F. Strniste, and H. Nagasawa, Sublethal damage repair and G2 block of xrs-6 cells to X-irradiation, manuscript in preparation (1991)
32. J. Y. Chan, and D. J. Chen, DNA Ligase Activities in a Radiation-Sensitive Mutant XRS-6 Cells, *Mutat. Res.*, Submitted (1992)
33. Warters, R. L., M. T. Li, B. W. Lyons, and D. J. Chen, Topoisomerase II Activity in a DNA Double-Strand Break Repair Mutant, *Radiat. Res.* 254:167-174 (1991)
34. R. A. Toby and K. D. Ley, Isoleucine-mediated regulation of genome replication in various mammalian cell lines, *Cancer Res.* 31:46-51 (1971)
35. P. E. Bryant, Enzymatic restriction of mammalian cells using Pvu-II and Bam HI: evidence for the double-strand break origin of chromosomal aberrations, *Int. J. Radiat. Biol.* 46:57-65 (1984)
36. H. H. Evans, J. Mencl, M. F. Horng, M. Ricanati, C. Sanchez and J. Hozier, Locus specificity in the mutability of L5178Y mouse lymphoma cells: The role of mutilocus lesions, *Proc. Natl. Acad. Sci. (USA)*, 83:4379-4383 (1986)
37. S. Tadahiro, N. Hieda-Shiomi, K. Sato, H. Tsuji, E. Takahashi and I. Tobari, A mouse-cell mutant sensitive to ionizing radiation is hypermutable by low dose of γ -radiation, *Mutat. Res.* 83:107-116 (1981)
38. I. Furuno Fukushima and H. Matsudaira, Mutation induction by different dose rates of γ -rays in radiation-sensitive mutants of mouse leukemia cells, *Radiat. Res.* 120:370-374 (1989)
39. P. A. Jeggo, M. Hafezparast, A. F. Thompson, B. C. Broughton, G. P. Kaur, M. Z. Zdzienicka and r. S. Athwal, Olcatization of a DNA repair gene (XRCC%) involved in double strand break rejoining to human chromosome 2, *Proc. Natl. Acad. Sci. (USA)* 1992
40. R. A. Gibbs, J. Camakaris, G. S. Hodgson and R. F. Martin, Molecular characterization of 125I decay and X-ray induced HPRT mutants in CHO cells, *Int. J. Radiat. Biol.* 51:193-199 (1987)
41. J. W. Szostak, T. E. Orr-Weaver, R. L. Rothstein and F. W. Stahl, The double-strand break repair model for combination, *Cell* 33:25-35 (1983)
42. A. Westerveld, J.H.H. Hoeijmakers, M. van Duin, J. de Wit, H. Odijk, A. Pastink, R.D. Wood and D. Bootsma, Molecular cloning of a human DNA repair gene, *Nature* 310:425-429 (1984)
43. C.A. Weber, D. P. Salazar, S.A. Stewart and L.H. Thompson, Molecular cloning and biological characterization of a human gene ERCC5, that corrects the nucleotide excision repair deficient in CHO UV35 cells, *Mol. Cell Biol.* 8:1137-1146 (1985)
44. J. S. Mudgett and M. A. MacInnes, Isolation of the functional human excision repair gene ERCC5 by intercosmid recombination, *Genomics* 8:623-633 (1990)
45. G. Weeda, R. C.A. van Ham, R. Masurel, A. Westerveld, H. Odijk, J. de Wit, D. Bootsma, A. van der EB andf J H. J. Hoeijmakers, Molecular cloning and biological characterization of the human excision repair gene ERCC3, *Mole. Cell. Biol.* 10:2570-2581 (1990)
46. L. H. Thompson, K. W. Brookman, N. J. Jones, S. A. Allen, and A. V. Carrano, Molecular cloning of the human XRCC1 gene, which corrects defective DNA strand-break repair and sister chromatid exchange, *Mole. Cell Biol.* in press (1991)
47. Siciliano, M.J. , A.V. Carrano and L.H. Thompson, Assignment of a human DNA-repair gene associated with sister-chromatid exchange to chromosome 19, *Mutat. Res.* 174:303-308 (1986)
48. M. Koi, M. Shimizu, H. Morita, H. Yamada and M. Oshimura, Construction of mouse A9 clones containing a single human chromosome tagged with neomycin-resistance gene via microcell fusion, *Jpn. J. Cancer Res.* 80:413-418 (1989)
49. L. H. Thompson, A. V. Carrano, K. Sato, E. P. Salazar, b. f. White, S. A. Stewart, J. L. Minkler, and M. J. Siciliano, Identification of nucleotide-excision-repair genes

- on human chromosomes 2 and 13 by functional complementation in hamster-human hybrids, *Somat. Cell Mol. Genet.* 13:539-551 (1987)
50. R.S. Atheal, M. Smarsh, B. Searle, and S. S. Deo, Integration of a dominant selectable marker into human chromosomes and transfer of marked chromosomes to mouse cells by microcell fusion, *Somat. Cell Genet.*, 11:177-187 (1985)
 51. T. G. Lugo, B. Handelin, A. M. Killary, D. E. Housman, Isolation of Microcell hybrid clones containing retroviral vector insertions into specific human chromosomes, *Mol. Cell. Biol.* 7:2814-2820 (1987)
 52. D. R. Cox, C. A. Pritchard, E. Uglum, D. Casher, I. Kobor, and R. M. Myers, Segregation of the huntington disease region of human chromosome 4 in a somatic cell hybrid, *Genomic* 4:397-407 (1989)
 53. D. L. Nelson, S. A. Ledbetter, L. Corbo, M. F. Victoria, R. Ramirez-Solis, T. D. Webster, D. H. Ledbetter and C.T. Caskey, Alu polymerase chain reaction: A method for rapid isolation of human-specific sequences from complex DNA sources, *Proc. Natl Acad. Sci. USA* 86:6686-6690 (1989)
 54. L. H. Thompson, Mooney, C. L., Brukhardt-Schultz, K., Carrano, A. V. and Siciliano, M. J. (1985). Correction of a nucleotide-excision-repair mutation by human chromosome 19 in hamster-human hybrid cells. *Somat. Cell Mol. Genet.* 11:87-92.
 55. M. J. Siciliano, Carrano, A. V. and Thompson, L. H. (1986). Assignment of a human DNA-repair gene associated with sister-chromatid
 56. L. H. Thompson, Bachinski, L., Weber, C. A., Stallings, R. Siciliano, M. J. (1989). Complementation of repair gene mutations on the hemizygous chromosome 9 in CHO cells: a third DNA repair gene on human chromosome 19. *Genomics* 5:670-679.
 57. A. J. Giaccia, Denko, N., MacLaren, R., Mirman, D., Waldren, C., Hart, I., and Stamato, T. D. (1990). Human chromosome 5 complements the DNA double-strand break-repair deficiency and gamma-ray sensitivity of the XR-1 hamster variant. *Am. J. Hum. Genet.* 47:359-469.
 58. P. Liu, Callen, C. F. Reeders, S. T., Sutherland, G. R., Thompson, L., Siciliano, M. J. (1989). Human DNA excision repair gene ERCC4 is located on chromosome 16 short arm 16p13-p13.3. *Cytogenet. Cell Genet.* 51:1035 (abstract).
 59. L. H. Thompson, Brookman, K. W., and Siciliano, M. J. (1991). Mammalian genes involved in the repair of damage from ionizing radiation, in *Ionizing Radiation Damage to DNA: Molecular Approaches*. UCLA Symposium Publication, in press.
 60. D. J. Chen, M. S. Park, E. Campbell, M. Oshimura, P. Liu, Y. Zhao, B. White and M. J. Siciliano, Assignment of a human DNA double-strand break repair gene (XRCC5) to chromosome 2, *Genomics* 13:1088-1094 (1992)
 61. K. Sato, A. Ito, T. Shiomi, H. Hama-Inaba, H. Ishikawa, T. Yoshizumi and T. Nakazawa, X-ray-sensitive mutants of mouse mammary carcinoma cells are hypersensitive to bleomycin and hydrogen peroxide, *Jpn. J. Cancer Res.* 77:456-461 (1986)
 62. G. F. Whitmore, A. J. Varghese and S. Gulyas, Cell cycle repair of double-strand DNA breaks in a gamma-ray-sensitive Chinese hamster cell, *Somat. Cell Mol. Genet.* 11:485-491 (1985)
 63. M. Z. Zdzienicka, N. van Welkel and G. P. van der Schana, A fourth complementation group among ionizing radiation-sensitive Chinese hamster cell mutant defective in DNA double-strand break repair, *Radiat. Res.* 131 (1992)
 64. C. N. Robson, A. Hall, A. L. Harris and I. D. Hickson, Bleomycin and X-ray-hypersensitive Chinese hamster ovary cells mutants: Genetic analysis and cross-resistance to neocarzinostatin, *Mutat. Res.* 193:157-165 (1988)
 65. R. L. Warters, B. W. Lyons, D. J. Chen and K. Sato, DNA damage processing in a radiation sensitive mouse cell line, *Mutat. Res.* in press (1992)

66. H. Nagasawa, A. B. Cox and J. T. Lett, The radiation responses of synchronous L5178Y S/S cells and thier significance for radiological theory, Proc. R. Soc. Lond. B211: 25-49 (1980)